



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$Q_c(a, \beta)$ does not contain p^2 , as we have just seen. But these two numbers contain no other factor in common, as is seen from Theorem I. Hence $Q_c(a, \beta)$ contains a prime not in $a^{c/p} - \beta^{c/p}$ at least in every case for which $Q_c(a, \beta) \neq p$ and also in every case for which $a^{c/p} - \beta^{c/p}$ is not divisible by p , unless $Q_c(a, \beta) = 1$.

COROLLARY. $a^c - \beta^c$ has always a prime factor not dividing $a^s - \beta^s$ ($s < c$) except in the cases mentioned in Theorem VII.

NOTE ON THE EXTENSION OF THE EXPONENTIAL THEOREM.

By E. D. ROE, JR., Syracuse University.

In the writer's paper in the June-July number of the MONTHLY, pp. 101-106, it will be observed that the complex roots, infinite in number, arising from the application of an incommensurable exponent are tacitly neglected, only the single real root being used. In fact this is the only root that is of much practical value. It is the only root that is usually considered in the extension of the binomial theorem for the expansion of $(1+x)^n$, even for a commensurable fractional exponent. The finite number of complex roots can be easily expressed if wanted. But when the exponent is incommensurable the number of complex roots becomes infinite and the complex roots become indeterminate.

In both cases, vis., of the binomial theorem, and the exponential theorem, which depends on the binomial theorem, the developments already obtained would have to be multiplied by $\cos \varphi + i \sin \varphi$, to obtain the complete development, where φ admits the value of zero, and has besides an infinite number of values which are indeterminate, when the exponent is incommensurable. All the values would have the same modulus.

These facts are so obvious that it will doubtless appear superfluous to many readers to call attention to them; yet for others it may not be amiss to do so.